

SKIN SERIOUSLY.

Cosmetic Fillers:

Restylane, Juvederm, Radiesse

Before Treatment:

- Avoid taking aspirin, non-steroidal anti-inflammatory medications, St. John's wort, and high doses of Vitamin E supplements for one week before and after treatment. These agents may increase bruising and bleeding at the injections site.
- If you have previously suffered from facial cold sores, there is a risk that the needle punctures could contribute to another occurrence. Speak to your physician about medications that may minimize a recurrence.

After Treatment:

- Cold compresses (a cloth dipped in cold water, wrung out, and applied to the injected area) or ice pack may be used immediately after the treatment to reduce swelling.
- Avoid touching the treated area within six hours following the treatment so you do not accidentally injure your skin while the area is numb. If there are any nodules you can gently massage the area 24hrs after the treatment until they resolve (usually 1 week).
- Until there is no redness or swelling, avoid exposure of the treated area to intense heat (sun lamp or sun bathing).

Adrienne Rencic, MD, PhD
Crossroads Center
1102 Baltimore Pike
Suite 202
Glen Mills, PA 19342
Phone : 610.558.1446
Fax : 610.558.1449

Hours:

Monday:

12:00 pm - 7:00 pm

Tuesday - Friday:

8:00 am - 5:00 pm